

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification: <b>A61K 31/40, 31/44, 31/47, C07D 207/08, 207/09, 213/89, 215/60</b>	<b>A1</b>	(11) International Publication Number: <b>WO 98/37885</b> (43) International Publication Date: 3 September 1998 (03.09.98)
(1) International Application Number: <b>PCT/US98/03484</b> (22) International Filing Date: 26 February 1998 (26.02.98) (30) Priority Data: 08/807,406                      28 February 1997 (28.02.97)      US (71) Applicant: <b>GUILFORD PHARMACEUTICALS INC.</b> [US/US]; 6611 Tributary Street, Baltimore, MD 21224 (US). (72) Inventors: <b>HAMILTON, Gregory, S.</b> ; 6501 Frederick Road, Catonsville, MD 21228 (US). <b>STEINER, Joseph, P.</b> ; 988 Sugar Maple Street, Hampstead, MD 21074 (US). <b>BURAK, Eric, S.</b> ; 3004 Victoria's Way, Forest Hill, MD 21050 (US). (74) Agent: <b>NATH, Gary, M.</b> ; Nath & Associates, Suite 750, 1835 K Street, N.W., Washington, DC 20006-1203 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>

(4) Title: **N-OXIDES OF HETEROCYCLIC ESTERS, AMIDES, THIOESTERS, AND KETONES**

(57) Abstract

This invention relates to neurotrophic low molecular weight, small molecule N-oxides of heterocyclic esters, amides, thioesters, and ketones having an affinity for FKBP-type immunophilins, and their use as inhibitors of the enzyme activity associated with immunophilin proteins, particularly peptidyl-prolyl isomerase, or rotamase, enzyme activity.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

N-OXIDES OF HETEROCYCLIC ESTERS, AMIDES, THIOESTERS,  
AND KETONES

BACKGROUND OF THE INVENTION

5

1. Field of Invention

This invention relates to neurotrophic low molecular weight, small molecule N-oxides of heterocyclic esters having an affinity for FKBP-type immunophilins, and their use as inhibitors of the enzyme activity associated with immunophilin proteins, particularly peptidyl-prolyl isomerase, or rotamase, enzyme activity.

2. Description of Related Art

The term immunophilin refers to a number of proteins that serve as receptors for the principal immunosuppressant drugs, cyclosporin A (CsA), FK506 and rapamycin. Known classes of immunophilins are cyclophilins and FK506 binding proteins, or FKBP's. Cyclosporin A binds to cyclophilin A while FK506 and rapamycin bind to FKBP12. These immunophilin-drug complexes interface with various intracellular signal transduction systems, especially the immune and nervous systems.

Immunophilins are known to have peptidyl-prolyl isomerase (PPIase), or rotamase, enzyme activity. It has been determined that rotamase enzyme activity plays a role in the catalyzation of the interconversion of the cis and trans isomers of peptide and protein substrates for the immunophilin proteins.

Immunophilins were originally discovered and

studied in the immune tissue. It was initially postulated by those skilled in the art that inhibition of the immunophilins' rotamase activity leads to inhibition of T-cell proliferation, thereby causing the immunosuppressive activity exhibited by immunosuppressant drugs, such as cyclosporin A, FK506 and rapamycin. Further study has shown that the inhibition of rotamase activity, in and of itself, does not result in immunosuppressive activity. Schreiber et al., *Science*, 1990, vol. 250, pp. 556-559. Instead, immunosuppression appears to stem from the formulation of a complex of immunosuppressant drugs and immunophilins. It has been shown that the immunophilin-drug complexes interact with ternary protein targets as their mode of action. Schreiber et al., *Cell*, 1991, vol. 66, pp. 807-815. In the case of FKBP-FK506 and cyclophilin-CsA, the immunophilin-drug complexes bind to the enzyme calcineurin and inhibit the T-cell receptor signalling which leads to T-cell proliferation. Similarly, the immunophilin-drug complex of FKBP-rapamycin interacts with the RAFT1/FRAP protein and inhibits the IL-2 receptor signalling.

Immunophilins have been found to be present at high concentrations in the central nervous system. Immunophilins are enriched 10-50 times more in the central nervous system than in the immune system. Within neural tissues, immunophilins appear to

influence nitric oxide synthesis, neurotransmitter release and neuronal process extension.

It has been found that picomolar concentrations of an immunosuppressant such as FK506 and rapamycin stimulate neurite outgrowth in PC12 cells and sensory neurons, namely dorsal root ganglion cells (DRGs). Lyons et al., *Proc. of Natl. Acad. Sci.*, 1994, vol. 91, pp. 3191-3195. In whole animal experiments, FK506 has been shown to stimulate nerve regeneration following facial nerve injury.

Surprisingly, it has been found that certain compounds with a high affinity for FKBP are potent rotamase inhibitors and exhibit excellent neurotrophic effects. Furthermore, these rotamase inhibitors are devoid of immunosuppressive activity. These findings suggest the use of rotamase inhibitors in treating various peripheral neuropathies and enhancing neuronal regrowth in the central nervous system (CNS). Studies have demonstrated that neurodegenerative disorders such as Alzheimer's disease, Parkinson's disease, and amyotrophic lateral sclerosis (ALS) may occur due to the loss, or decreased availability, of a neurotrophic substance specific for a particular population of neurons affected in the disorder.

Several neurotrophic factors affecting specific neuronal populations in the central nervous system have been identified. For example, it has been hypothesized that Alzheimer's disease results from a

and neurotrophin 3,  
to increase the survival of degenerating neuronal  
populations.


Clinical application of these proteins in various  
10 neurological disease states is hampered by  
difficulties in the delivery and bioavailability of  
large proteins to nervous system targets. By  
contrast, immunosuppressant drugs with neurotrophic  
activity are relatively small and display excellent  
15 bioavailability and specificity. However, when  
administered chronically, immunosuppressant drugs  
exhibit a number of potentially serious side effects  
including nephrotoxicity, such as impairment of  
glomerular filtration and irreversible interstitial  
20 fibrosis (Kopp et al., *J. Am. Soc. Nephrol.*, 1991,  
1:162); neurological deficits, such as involuntary  
tremors, or non-specific cerebral angina, such as non-  
localized headaches (De Groen et al., *N. Engl. J.*  
*Med.*, 1987, 317:861); and vascular hypertension with  
25 complications resulting therefrom (Kahan et al., *N.*  
*Engl. J. Med.*, 1989, 321:1725).

In order to prevent the side effects associated  
with use of the immunosuppressant compounds, the



compared to compounds lacking the N-oxide group.

Specifically, the present invention relates to a compound of formula I:



I

or a pharmaceutically acceptable salt thereof, wherein:

A and B are taken together, with the nitrogen and carbon atoms to which they are respectively attached, to form a 5-7 membered saturated or unsaturated heterocyclic ring containing any combination of CH<sub>2</sub>, O, S, SO, SO<sub>2</sub>, NH or NR<sub>1</sub> in any chemically stable oxidation state;

W is O, S, CH<sub>2</sub>, or H<sub>2</sub>;

R is a C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or alkenyl group optionally substituted with C<sub>3</sub>-C<sub>8</sub> cycloalkyl, C<sub>3</sub> or C<sub>5</sub> cycloalkyl, C<sub>5</sub>-C<sub>7</sub> cycloalkenyl, or Ar<sub>1</sub>, where said alkyl, alkenyl, cycloalkyl, or cycloalkenyl groups may be optionally substituted with C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkenyl, or hydroxy, and where Ar<sub>1</sub> is selected from the group consisting of 1-naphthyl, 2-naphthyl, 1-indolyl, 2-indolyl, 2-furyl, 3-furyl, 2-



thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, or phenyl, having one to three substituents which are independently selected from the group consisting of hydrogen, halo, hydroxyl, nitro, trifluoromethyl, C<sub>1</sub>-C<sub>6</sub> straight or branched alkyl or alkenyl, C<sub>1</sub>-C<sub>4</sub> alkenyloxy, phenoxy, benzyloxy, and amino;

X is O, NH, NR<sub>1</sub>, S, CH, CR<sub>1</sub>, or C(R<sub>1</sub>)<sub>2</sub>;


Y is a direct bond, or a C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or alkenyl which is optionally substituted in one or more positions with C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or alkenyl, or C<sub>3</sub>-C<sub>8</sub> cycloalkyl, or C<sub>3</sub>-C<sub>7</sub> cycloalkenyl, or hydroxyl, or carbonyl oxygen, or with Ar, where said alkyl, alkenyl, cycloalkyl, cycloalkenyl, or Ar group is optionally substituted with C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkenyl, or hydroxy, or carbonyl oxygen, or wherein any of the carbon atoms of said alkyl, alkenyl, cycloalkyl, cycloalkenyl, or Ar group is optionally replaced with O, NH, NR<sub>2</sub>, S, SO, or SO<sub>2</sub>, where R<sub>2</sub> is selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>4</sub>)-straight or branched chain alkyl, (C<sub>3</sub>-C<sub>4</sub>)-straight or branched chain alkenyl or alkynyl, and (C<sub>1</sub>-C<sub>4</sub>) bridging alkyl wherein a bridge is formed between the nitrogen and a carbon atom of said alkyl or alkenyl chain containing said heteroatom to form a ring, wherein said ring is optionally fused to an Ar group; and

Z is an aromatic or tertiary alkyl amine oxidized to a corresponding N-oxide, wherein the aromatic amine

oxidized to a corresponding N-oxide where Ar is  
a mono-, bi- or tricyclic, carbo- or heterocyclic  
ring, wherein the ring is either unsubstituted or  
substituted in one to three position(s) with halo,  
5 hydroxyl, nitro, trifluoromethyl, C<sub>1</sub>-C<sub>6</sub> straight or  
branched chain alkyl or alkenyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub>  
alkenyloxy, phenoxy, benzyloxy, amino, or a  
combination thereof; wherein the individual ring sizes  
are 5-6 members; wherein the heterocyclic ring  
10 contains 1-6 heteroatom(s) selected from the group  
consisting of O, N, S, and a combination thereof  
wherein at least one of the heteroatoms is N, and  
wherein the alkyl amine is oxidized to a corresponding  
N-oxide where alkyl is a C<sub>1</sub>-C<sub>6</sub> straight or branched  
15 chain alkyl or alkenyl which is optionally substituted  
in one or more positions with C<sub>1</sub>-C<sub>6</sub> straight or  
branched chain alkyl or alkenyl, or C<sub>3</sub>-C<sub>8</sub> cycloalkyl,  
or C<sub>5</sub>-C<sub>7</sub> cycloalkenyl, or hydroxyl, or carbonyl oxygen,  
or with Ar, where said alkyl, alkenyl, cycloalkyl,  
20 cycloalkenyl, or Ar group is optionally substituted  
with C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkenyl, or hydroxy, or carbonyl  
oxygen, or wherein any of the carbon atoms of said  
alkyl, alkenyl, cycloalkyl, cycloalkenyl, or Ar group  
is optionally replaced with O, NH, NR<sub>1</sub>, S, SO, or SO<sub>2</sub>;  
25 and,

R<sub>1</sub> is hydrogen, (C<sub>1</sub>-C<sub>4</sub>)-straight or branched chain  
alkyl, (C<sub>3</sub>-C<sub>4</sub>)-straight or branched chain alkenyl or  
alkynyl, or R<sub>1</sub> is Y-Z, as defined above.

Another preferred embodiment of this invention is a compound of formula II:



II

or a pharmaceutically acceptable salt thereof,  
wherein:

E, F, G and H are independently CH<sub>2</sub>, O, S, SO,  
SO<sub>2</sub>, NH or NR<sub>1</sub>;

W is O, S, CH<sub>2</sub>, or H<sub>2</sub>;

R is a C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or alkenyl group optionally substituted with C<sub>3</sub>-C<sub>8</sub> cycloalkyl, C<sub>3</sub> or C<sub>5</sub> cycloalkyl, C<sub>5</sub>-C<sub>7</sub> cycloalkenyl, or Ar<sub>1</sub>, where said alkyl, alkenyl, cycloalkyl, or cycloalkenyl groups may be optionally substituted with C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkenyl, or hydroxy, and where Ar<sub>1</sub> is selected from the group consisting of 1-naphthyl, 2-naphthyl, 1-indolyl, 2-indolyl, 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, or phenyl, having one to three substituents which are independently selected from the group consisting of hydrogen, halo, hydroxyl, nitro, trifluoromethyl, C<sub>1</sub>-C<sub>6</sub> straight or branched alkyl or alkenyl, C<sub>1</sub>-C<sub>4</sub> alkenyloxy, phenoxy, benzyloxy, and amino;

20


substituted in one or more positions with C<sub>1</sub>-C<sub>6</sub>  
5 straight or branched chain alkyl or alkenyl, or C<sub>3</sub>-C<sub>8</sub>  
cycloalkyl, or C<sub>5</sub>-C<sub>8</sub> cycloalkenyl, or hydroxyl, or  
carbonyl oxygen, or with Ar, where said alkyl,  
alkenyl, cycloalkyl, cycloalkenyl, or Ar group is  
optionally substituted with C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkenyl,  
10 or hydroxy, or carbonyl oxygen, or wherein any of the  
carbon atoms of said alkyl, alkenyl, cycloalkyl,  
cycloalkenyl, or Ar group is optionally replaced with  
O, NH, NR<sub>2</sub>, S, SO, or SO<sub>2</sub>, where R<sub>2</sub> is selected from  
the group consisting of hydrogen, (C<sub>1</sub>-C<sub>4</sub>)-straight or  
15 branched chain alkyl, (C<sub>3</sub>-C<sub>4</sub>)-straight or branched  
chain alkenyl or alkynyl, and (C<sub>1</sub>-C<sub>4</sub>) bridging alkyl  
wherein a bridge is formed between the nitrogen and a  
carbon atom of said alkyl or alkenyl chain containing  
said heteroatom to form a ring, wherein said ring is  
20 optionally fused to an Ar group; and

Z is an aromatic or tertiary alkyl amine oxidized  
to a corresponding N-oxide, wherein the aromatic amine  
is Ar oxidized to a corresponding N-oxide where Ar is  
a mono-, bi- or tricyclic, carbo- or heterocyclic  
25 ring, wherein the ring is either unsubstituted or  
substituted in one to three position(s) with halo,  
hydroxyl, nitro, trifluoromethyl, C<sub>1</sub>-C<sub>6</sub> straight or  
branched chain alkyl or alkenyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub>

alkenyloxy, phenoxy, benzyloxy, amino, or a combination thereof; wherein the individual ring sizes are 5-6 members; wherein the heterocyclic ring contains 1-6 heteroatom(s) selected from the group consisting of O, N, S, and a combination thereof wherein at least one of the heteroatoms is N, and wherein the alkyl amine is oxidized to a corresponding N-oxide where alkyl is a C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or alkenyl which is optionally substituted in one or more positions with C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or alkenyl, or C<sub>3</sub>-C<sub>8</sub> cycloalkyl, or C<sub>5</sub>-C<sub>7</sub> cycloalkenyl, or hydroxyl, or carbonyl oxygen, or with Ar, where said alkyl, alkenyl, cycloalkyl, cycloalkenyl, or Ar group is optionally substituted with C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkenyl, or hydroxy, or carbonyl oxygen, or wherein any of the carbon atoms of said alkyl, alkenyl, cycloalkyl, cycloalkenyl, or Ar group is optionally replaced with O, NH, NR<sub>1</sub>, S, SO, or SO<sub>2</sub>; and,

R<sub>1</sub> is hydrogen, (C<sub>1</sub>-C<sub>4</sub>)-straight or branched chain alkyl, (C<sub>3</sub>-C<sub>4</sub>)-straight or branched chain alkenyl or alkynyl, or R<sub>1</sub> is Y-Z, as defined above.

Another preferred embodiment is a compound of formula III:



III

or a pharmaceutically acceptable salt thereof,  
 5 wherein:

E, F, and G are independently  $\text{CH}_2$ , O, S, SO,  $\text{SO}_2$ ,  
 NH or  $\text{NR}_1$ ;

W is O, S,  $\text{CH}_2$ , or  $\text{H}_2$ ;

R is a  $\text{C}_1$ - $\text{C}_6$  straight or branched chain alkyl or  
 10 alkenyl group optionally substituted with  $\text{C}_1$ - $\text{C}_6$   
 cycloalkyl,  $\text{C}_3$  or  $\text{C}_5$  cycloalkyl,  $\text{C}_5$ - $\text{C}_7$  cycloalkenyl, or  
 $\text{Ar}_1$ , where said alkyl, alkenyl, cycloalkyl, or  
 cycloalkenyl groups may be optionally substituted with  
 $\text{C}_1$ - $\text{C}_4$  alkyl,  $\text{C}_1$ - $\text{C}_4$  alkenyl, or hydroxy, and where  $\text{Ar}_1$  is  
 15 selected from the group consisting of 1-naphthyl, 2-  
 naphthyl, 1-indolyl, 2-indolyl, 2-furyl, 3-furyl, 2-  
 thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl,  
 or phenyl, having one to three substituents which are  
 independently selected from the group consisting of  
 20 hydrogen, halo, hydroxyl, nitro, trifluoromethyl,  $\text{C}_1$ - $\text{C}_6$   
 straight or branched alkyl or alkenyl,  $\text{C}_1$ - $\text{C}_4$   
 alkenyloxy, phenoxy, benzyloxy, and amino;

X is O, NH,  $\text{NR}_1$ , S, CH,  $\text{CR}_1$ , or  $\text{C}(\text{R}_1)_2$ ;

Y is a direct bond, or a  $\text{C}_1$ - $\text{C}_6$  straight or  
 25 branched chain alkyl or alkenyl which is optionally


substituted in one or more positions with C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or alkenyl, or C<sub>3</sub>-C<sub>8</sub> cycloalkyl, or C<sub>5</sub>-C<sub>7</sub> cycloalkenyl, or hydroxyl, or carbonyl oxygen, or with Ar, where said alkyl, alkenyl, cycloalkyl, cycloalkenyl, or Ar group is optionally substituted with C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkenyl, or hydroxy, or carbonyl oxygen, or wherein any of the carbon atoms of said alkyl, alkenyl, cycloalkyl, cycloalkenyl, or Ar group is optionally replaced with O, NH, NR<sub>2</sub>, S, SO, or SO<sub>2</sub>, where R<sub>2</sub> is selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>4</sub>)-straight or branched chain alkyl, (C<sub>3</sub>-C<sub>4</sub>)-straight or branched chain alkenyl or alkynyl, and (C<sub>1</sub>-C<sub>4</sub>) bridging alkyl wherein a bridge is formed between the nitrogen and a carbon atom of said alkyl or alkenyl chain containing said heteroatom to form a ring, wherein said ring is optionally fused to an Ar group; and

Z is an aromatic or tertiary alkyl amine oxidized to a corresponding N-oxide, wherein the aromatic amine is Ar oxidized to a corresponding N-oxide where Ar is a mono-, bi- or tricyclic, carbo- or heterocyclic ring, wherein the ring is either unsubstituted or substituted in one to three position(s) with halo, hydroxyl, nitro, trifluoromethyl, C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or alkenyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> alkenyloxy, phenoxy, benzyloxy, amino, or a combination thereof; wherein the individual ring sizes are 5-6 members; wherein the heterocyclic ring

contains 1-6 heteroatom(s) selected from the group consisting of O, N, S, and a combination thereof wherein at least one of the heteroatoms is N, and wherein the alkyl amine is oxidized to a corresponding N-oxide where alkyl is a C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or alkenyl which is optionally substituted in one or more positions with C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or alkenyl, or C<sub>3</sub>-C<sub>8</sub> cycloalkyl, or C<sub>5</sub>-C<sub>7</sub> cycloalkenyl, or hydroxyl, or carbonyl oxygen, or with Ar, where said alkyl, alkenyl, cycloalkyl, cycloalkenyl, or Ar group is optionally substituted with C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkenyl, or hydroxy, or carbonyl oxygen, or wherein any of the carbon atoms of said alkyl, alkenyl, cycloalkyl, cycloalkenyl, or Ar group is optionally replaced with O, NH, NR<sub>1</sub>, S, SO, or SO<sub>2</sub>; and,

R<sub>1</sub> is hydrogen, (C<sub>1</sub>-C<sub>4</sub>)-straight or branched chain alkyl, (C<sub>3</sub>-C<sub>4</sub>)-straight or branched chain alkenyl or alkynyl, or R<sub>1</sub> is Y-Z, as defined above.

A further particularly preferred embodiment of this invention is a compound of formula IV:



IV

or a pharmaceutically acceptable salt thereof,



wherein:

n is 1, 2 or 3 forming a 5-7 member heterocyclic ring;

W is O, S, CH<sub>2</sub>, or H<sub>2</sub>;

5 R is a C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or alkenyl group optionally substituted with C<sub>3</sub>-C<sub>8</sub> cycloalkyl, C<sub>3</sub> or C<sub>5</sub> cycloalkyl, C<sub>5</sub>-C<sub>7</sub> cycloalkenyl, or Ar<sub>1</sub>, where said alkyl, alkenyl, cycloalkyl, or cycloalkenyl groups may be optionally substituted with  
10 C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkenyl, or hydroxy, and where Ar<sub>1</sub> is selected from the group consisting of 1-naphthyl, 2-naphthyl, 1-indolyl, 2-indolyl, 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, or phenyl, having one to three substituents which are  
15 independently selected from the group consisting of hydrogen, halo, hydroxyl, nitro, trifluoromethyl, C<sub>1</sub>-C<sub>6</sub> straight or branched alkyl or alkenyl, C<sub>1</sub>-C<sub>4</sub> alkenyloxy, phenoxy, benzyloxy, and amino;

X is O, NH, NR<sub>1</sub>, S, CH, CR<sub>1</sub>, or C(R<sub>1</sub>)<sub>2</sub>;

20 Y is a direct bond, or a C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or alkenyl which is optionally substituted in one or more positions with C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or alkenyl, or C<sub>3</sub>-C<sub>8</sub> cycloalkyl, or C<sub>5</sub>-C<sub>7</sub> cycloalkenyl, or hydroxyl, or  
25 carbonyl oxygen, or with Ar, where said alkyl, alkenyl, cycloalkyl, cycloalkenyl, or Ar group is optionally substituted with C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkenyl, or hydroxy, or carbonyl oxygen, or wherein any of the

carbon atoms of said alkyl, alkenyl, cycloalkyl, cycloalkenyl, or Ar group is optionally replaced with O, NH, NR<sub>2</sub>, S, SO, or SO<sub>2</sub>, where R<sub>2</sub> is selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>4</sub>)-straight or  
5 branched chain alkyl, (C<sub>3</sub>-C<sub>4</sub>)-straight or branched chain alkenyl or alkynyl, and (C<sub>1</sub>-C<sub>4</sub>) bridging alkyl wherein a bridge is formed between the nitrogen and a carbon atom of said alkyl or alkenyl chain containing said heteroatom to form a ring, wherein said ring is  
10 optionally fused to an Ar group; and

Z is an aromatic or tertiary alkyl amine oxidized to a corresponding N-oxide, wherein the aromatic amine is Ar oxidized to a corresponding N-oxide where Ar is a mono-, bi- or tricyclic, carbo- or heterocyclic  
15 ring, wherein the ring is either unsubstituted or substituted in one to three position(s) with halo, hydroxyl, nitro, trifluoromethyl, C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or alkenyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> alkenyloxy, phenoxy, benzyloxy, amino, or a  
20 combination thereof; wherein the individual ring sizes are 5-6 members; wherein the heterocyclic ring contains 1-6 heteroatom(s) selected from the group consisting of O, N, S, and a combination thereof wherein at least one of the heteroatoms is N, and  
25 wherein the alkyl amine is oxidized to a corresponding N-oxide where alkyl is a C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or alkenyl which is optionally substituted in one or more positions with C<sub>1</sub>-C<sub>6</sub> straight or

branched chain alkyl or alkenyl, or C<sub>3</sub>-C<sub>8</sub> cycloalkyl, or C<sub>5</sub>-C<sub>7</sub> cycloalkenyl, or hydroxyl, or carbonyl oxygen, or with Ar, where said alkyl, alkenyl, cycloalkyl, cycloalkenyl, or Ar group is optionally substituted with C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkenyl, or hydroxy, or carbonyl oxygen, or wherein any of the carbon atoms of said alkyl, alkenyl, cycloalkyl, cycloalkenyl, or Ar group is optionally replaced with O, NH, NR<sub>1</sub>, S, SO, or SO<sub>2</sub>; and,

R<sub>1</sub> is hydrogen, (C<sub>1</sub>-C<sub>4</sub>)-straight or branched chain alkyl, (C<sub>3</sub>-C<sub>4</sub>)-straight or branched chain alkenyl or alkynyl, or R<sub>1</sub> is Y-Z, as defined above.

In preferred embodiments, Ar is selected from the group consisting of pyrrolidinyl, pyridyl, pyrimidyl, pyrazyl, pyridazyl, quinolinyl, and isoquinolinyl.

Particularly preferred compounds of the present invention include:

3-(2-Pyridyl)-1-propyl (2S)-1-(3,3-Dimethyl-1,2-dioxo-pentyl)-2-pyrrolidinecarboxylate, N-oxide;

3-(3-Pyridyl)-1-propyl (2S)-1-(3,3-Dimethyl-1,2-dioxo-pentyl)-2-pyrrolidinecarboxylate, N-oxide;

3-(4-Pyridyl)-1-propyl (2S)-1-(3,3-Dimethyl-1,2-dioxo-pentyl)-2-pyrrolidinecarboxylate, N-oxide;

3-(2-Quinolyl)-1-propyl (2S)-1-(1,1-Dimethyl-1,2-dioxo-pentyl)-2-pyrrolidinecarboxylate, N-oxide;

3-(3-Quinolyl)-1-propyl (2S)-1-(1,1-Dimethyl-1,2-dioxo-pentyl)-2-pyrrolidinecarboxylate, N-oxide; and

3-(4-Quinolyl)-1-propyl(2S)-1-(1,1-Dimethyl-1,2-dioxo-pentyl)-2-pyrrolidinecarboxylate, N-oxide.

The present invention also relates to a pharmaceutical composition comprising a neurotrophically effective amount of the compound of formula I, II, III, or IV, and a pharmaceutically acceptable carrier.

The present invention further includes methods of using the compounds of the present invention. One preferred embodiment includes a method of stimulating damaged neurons in an animal, comprising:

administering to the animal a neurotrophically effective amount of a compound of the present invention.

Another preferred embodiment of the present invention includes a method of promoting neuronal regeneration in an animal, comprising:

administering to the animal a neurotrophically effective amount of a compound of the present invention.

Yet another embodiment of the present invention includes a method of preventing neurodegeneration in an animal, comprising:

administering to the animal a neurotrophically effective amount of a compound of the present invention.

Another embodiment includes a method of treating neurological disorders in an animal, comprising:

administering to the animal a neurotrophically effective amount of a compound of the present invention.

The neurological disorders for which the compounds of the present invention are particularly useful are selected from the group consisting of: peripheral neuropathy caused by physical injury or disease state, physical damage to the brain, physical damage to the spinal cord, stroke associated with brain damage, and neurological disorder relating to neurodegeneration. Examples of neurological disorders relating to neurodegeneration are Alzheimer's Disease, Parkinson's Disease, and amyotrophic lateral sclerosis.

#### BRIEF DESCRIPTION OF THE INVENTION

Figure 1 is a bar graph showing Protection of Striatal TH Innervation Density from MPTP-toxicity by Concurrent Drug Dosing. Figure 1 shows that at administration of 4 mg/kg of the compounds of the present invention there is a remarkable protection of striatal nerves from MPTP-toxicity.

#### DETAILED DESCRIPTION OF THE INVENTION

##### Definitions

"Alkyl" means a branched or unbranched saturated hydrocarbon chain containing 1 to 6 carbon atoms, such as methyl, ethyl, propyl, iso-propyl, butyl, iso-

butyl, tert-butyl, n-pentyl, n-hexyl, and the like, unless otherwise indicated.

"Alkoxy" means the group -OR wherein R is alkyl as herein defined. Preferably, R is a branched or  
5 unbranched saturated hydrocarbon chain containing 1 to 3 carbon atoms.

"Halo" means fluoro, chloro, bromo, or iodo, unless otherwise indicated.

"Phenyl" includes all possible isomeric phenyl  
10 radicals, optionally monosubstituted or multi-substituted with substituents selected from the group consisting of alkyl, alkoxy, hydroxy, halo, and haloalkyl.

The term "C<sub>1</sub>-C<sub>6</sub>" and similar terminology found in  
15 standard chemical nomenclature, when used for alkyl and alkenyl chains, is known in the art to include subchains such as C<sub>1</sub>-C<sub>3</sub>, C<sub>1</sub>-C<sub>4</sub>, C<sub>1</sub>-C<sub>5</sub>, C<sub>1</sub>-C<sub>6</sub>, C<sub>2</sub>-C<sub>4</sub>, C<sub>2</sub>-C<sub>5</sub>, C<sub>2</sub>-C<sub>6</sub>, C<sub>3</sub>-C<sub>5</sub>, C<sub>3</sub>-C<sub>6</sub>, C<sub>4</sub>-C<sub>6</sub>, and the variants thereof.

The term "pharmaceutically acceptable salt"  
20 refers to salts of the subject compounds which possess the desired pharmacological activity and which are neither biologically nor otherwise undesirable. The salts can be formed with inorganic acids such as acetate, adipate, alginate, aspartate, benzoate,  
25 benzenesulfonate, bisulfate, butyrate, citrate, camphorate, camphorsulfonate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, fumarate, glucoheptanoate, glycerophosphate,

hemisulfate heptanoate, hexanoate, hydrochloride  
hydrobromide, hydroiodide, 2-hydroxyethanesulfonate,  
lactate, maleate, methanesulfonate, 2-  
naphthalenesulfonate, nicotinate, oxalate, thiocya-  
5 nate, tosylate and undecanoate. Base salts include  
ammonium salts, alkali metal salts such as sodium and  
potassium salts, alkaline earth metal salts such as  
calcium and magnesium salts, salt with organic bases  
such as dicyclohexylamine salts, N-methyl-D-glucamine,  
10 and salts with amino acids such as arginine, lysine,  
and so forth. Also, the basic nitrogen-containing  
groups can be quarternized with such agents as lower  
alkyl halides, such as methyl, ethyl, propyl, and  
butyl chloride, bromides and iodides; dialkyl sulfates  
15 like dimethyl, diethyl, dibutyl and diamyl sulfates,  
long chain halides such as decyl, lauryl, myristyl and  
stearyl chlorides, bromides and iodides, aralkyl  
halides like benzyl and phenethyl bromides and others.  
Water or oil-soluble or dispersible products are  
20 thereby obtained.

The compounds of this invention possess at least  
one asymmetric center and thus can be produced as  
mixtures of stereoisomers or as individual enantiomers  
or diastereomers. The individual stereoisomers may be  
25 obtained by using an optically active starting  
material, by resolving a racemic or non-racemic  
mixture of an intermediate at some appropriate stage  
of the synthesis, or by resolution of the compound of

formula (I). It is understood that the individual stereoisomers as well as mixtures (racemic and non-racemic) of stereoisomers are encompassed by the scope of the present invention. The S-stereoisomer at atom 5 1 of formula I is most preferred due to its greater activity.

"Isomers" are different compounds that have the same molecular formula.

10 "Stereoisomers" are isomers that differ only in the way the atoms are arranged in space.

"Enantiomers" are a pair of stereoisomers that are non-superimposable mirror images of each other.

"Diastereoisomers" are stereoisomers which are not mirror images of each other.

15 "Racemic mixture" means a mixture containing equal parts of individual enantiomers. "Non-racemic mixture" is a mixture containing unequal parts of individual enantiomers or stereoisomers.

20 The term "treatment" as used herein covers any treatment of a disease and/or condition in an animal, particularly a human, and includes:

(i) preventing a disease and/or condition from occurring in a subject which may be predisposed to the disease and/or condition but has not yet been diagnosed as having it;

25

(ii) inhibiting the disease and/or condition, i.e., arresting its development; or

(iii) relieving the disease and/or condition,



i.e., causing regression of the disease and/or condition.

The term "effecting" as directed to "effecting neuronal activity" relates to producing or bringing about a desired effect or result and includes without limitation stimulating neurons, promoting regeneration preventing degeneration, protecting from degeneration and treating disorders.


The system used in naming the compounds of the present invention is shown below, using a compound of formula IV as an example.

A compound of the present invention, especially Formula IV, wherein n is 1, R is 1,1-dimethylpentyl, X is O, Y is  $(CH_2)_3$ , and Z is 3-pyridyl-N-oxide, is named 3-(3-Pyridyl)-1-propyl (2S)-1-(3,3-Dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate, N-oxide.

#### Compounds of the Invention

The neurotrophic low molecular weight, small molecule FKBP inhibitor compounds of this invention have an affinity for FKBP-type immunophilins, such as FKBP12. When the neurotrophic compounds of this invention are bound to an FKBP-type immunophilin, they have been found to inhibit the prolyl-peptidyl cis-trans isomerase activity, or rotamase, activity of the binding protein and unexpectedly stimulate neurite growth.

Specific exemplifications of these embodiments are presented in TABLE I.

TABLE I

No.	n	X	Y	Z	R
1	1	O	$(CH_2)_3$	3-Pyridyl N-oxide	1,1-dimethylpentyl
2	1	O	$(CH_2)_3$	2-Pyridyl N-oxide	1,1-dimethylpentyl
3	1	O	$(CH_2)_3$	4-Pyridyl N-oxide	1,1-dimethylpentyl
4	1	O	$(CH_2)_3$	2-Quinolyl N-oxide	1,1-dimethylpentyl
5	1	O	$(CH_2)_3$	3-Quinolyl N-oxide	1,1-dimethylpentyl
6	1	O	$(CH_2)_3$	4-Quinolyl N-oxide	1,1-dimethylpentyl

The most preferred compounds of formula IV are:

3-(2-Pyridyl)-1-propyl (2S)-1-(3,3-Dimethyl-1,2-dioxo-pentyl)-2-pyrrolidinecarboxylate, N-oxide;

3-(3-Pyridyl)-1-propyl (2S)-1-(3,3-Dimethyl-1,2-dioxo-pentyl)-2-pyrrolidinecarboxylate, N-oxide;

3-(4-Pyridyl)-1-propyl (2S)-1-(3,3-Dimethyl-1,2-dioxo-pentyl)-2-pyrrolidinecarboxylate, N-oxide;

3-(2-Quinolyl)-1-propyl (2S)-1-(1,1-Dimethyl-1,2-dioxo-pentyl)-2-pyrrolidinecarboxylate, N-oxide;

3-(3-Quinolyl)-1-propyl (2S)-1-(1,1-Dimethyl-1,2-dioxo-pentyl)-2-pyrrolidinecarboxylate, N-oxide; and

3-(4-Quinolyl)-1-propyl (2S)-1-(1,1-Dimethyl-1,2-dioxo-pentyl)-2-pyrrolidinecarboxylate, N-oxide.

The compounds of the present invention exist as stereoisomeric forms, either enantiomers or diastereoisomers. Included within the scope of the invention are the enantiomers, the racemic form, and diastereoisomeric mixtures. Enantiomers and diastereoisomers can be separated by methods known to those skilled in the art.

#### Methods of Using the Compounds of the Invention

The compounds of the present invention have an affinity for the FK506 binding protein, particularly FKBP12, which is present in the neuronal tissue. When the inventive compounds bind to FKBP in neuronal tissue, they exhibit excellent neurotrophic activity. This activity is useful in the stimulation of damaged

several neurological disorders known to be associated  
5 with neuronal degeneration and peripheral  
neuropathies.

For the foregoing reasons, the present invention  
further relates to a method of effecting a neuronal  
activity in an animal, comprising:

10 administering to the animal a neurotrophically  
effective amount of a compound of formula I, II, III  
or IV.

In a preferred embodiment, the neuronal activity  
is selected from the group consisting of stimulation  
15 of damaged neurons, promotion of neuronal  
regeneration, prevention of neurodegeneration,  
protection of neurodegeneration, and treatment of  
neurological disorder.

The neurological disorders that may be treated  
20 include but are not limited to: trigeminal neuralgia;  
glossopharyngeal neuralgia; Bell's Palsy; myasthenia  
gravis; muscular dystrophy; amyotrophic lateral  
sclerosis; progressive muscular atrophy; progressive  
bulbar inherited muscular atrophy; herniated; ruptured  
25 or prolapsed invertebrate disk syndromes; cervical  
spondylosis; plexus disorders; thoracic outlet  
destruction syndromes; peripheral neuropathic such as  
those caused by lead, dapsone, ticks, porphyria, or

Guillain-Barré syndrome; Alzheimer's disease; and Parkinson's disease.

5 The compounds of the present invention are particularly useful for treating a neurological disorder selected from the group consisting of: peripheral neuropathy caused by physical injury or disease state, physical damage to the brain, physical damage to the spinal cord, stroke associated with brain damage, and neurological disorder relating to  
10 neurodegeneration. Examples of neurological disorders relating to neurodegeneration are Alzheimer's Disease, Parkinson's Disease, and amyotrophic lateral sclerosis.

#### Pharmaceutical Compositions and Formulations

15 For these purposes the compounds of the present invention may be administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir in dosage formulations containing conventional non-toxic  
20 pharmaceutically-acceptable carriers, adjuvants and vehicles. The term parenteral as used herein includes subcutaneous, intravenous, intramuscular, intraperitoneally, intrathecally, intraventricularly, intrasternal and intracranial injection or infusion  
25 techniques.

To be effective therapeutically as central nervous system targets, the compounds of the present invention should readily penetrate the blood-brain barrier when

The compounds of the present invention may be administered in the form of sterile injectable preparations, for example, as sterile injectable aqueous or oleaginous suspensions. These suspensions  
10 may be formulated according to techniques known in the art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparations may also be sterile injectable solutions or suspensions in non-toxic parenterally-acceptable  
15 diluents or solvents, for example, as solutions in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally  
20 employed as solvents or suspending mediums. For this purpose, any bland fixed oil may be employed including synthetic mono- or di-glycerides. Fatty acids such as oleic acid and its glyceride derivatives, including olive oil and castor oil, especially in their  
25 polyoxyethylated versions, are useful in the preparation of injectables. These oil solutions or suspensions may also contain long-chain alcohol diluents or dispersants.

The compounds may be administered orally in the form of capsules, tablets, aqueous suspensions or solutions. Tablets may contain carriers such as lactose and corn starch, and/or lubricating agents such as magnesium stearate. Capsules may contain diluents including lactose and dried corn starch. Aqueous suspensions may contain emulsifying and suspending agents combined with the active ingredient. The oral dosage forms may further contain sweetening and/or flavoring and/or coloring agents.

The compounds of this invention may also be administered rectally in the form of suppositories. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at room temperature, but liquid at rectal temperature and, therefore, will melt in the rectum to release the drug. Such materials include cocoa butter, beeswax and polyethylene glycols.

The compounds of this invention may also be administered topically, especially when the conditions addressed for treatment involve areas or organs readily accessible by topical application, including neurological disorders of the eye, the skin, or the lower intestinal tract. Suitable topical formulations are readily prepared for each of these areas.

For topical application to the eye, or ophthalmic use, the compounds can be formulated as micronized suspensions in isotonic, pH adjusted sterile saline,

the ophthalmic uses the compounds may be formulated in  
5 an ointment such as petrolatum.

For topical application to the skin, the compounds  
can be formulated in a suitable ointment containing  
the compound suspended or dissolved in, for example,  
a mixture with one or more of the following: mineral  
10 oil, liquid petrolatum, white petrolatum, propylene  
glycol, polyoxyethylene polyoxypropylene compound,  
emulsifying wax and water. Alternatively, the  
compounds can be formulated in a suitable lotion or  
cream containing the active compound suspended or  
15 dissolved in, for example, a mixture of one or more of  
the following: mineral oil, sorbitan monostearate,  
polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-  
octyldodecanol, benzyl alcohol and water.

Topical application for the lower intestinal tract  
20 can be effected in a rectal suppository formulation  
(see above) or in a suitable enema formulation.

Dosage levels on the order of about 0.1 mg to  
about 10,000 mg of the active ingredient compound are  
useful in the treatment of the above conditions, with  
25 preferred levels of about 0.1 mg to about 1,000 mg.  
The amount of active ingredient that may be combined  
with the carrier materials to produce a single dosage  
form will vary depending upon the host treated and the



particular mode of administration.

It is understood, however, that a specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, rate of excretion, drug combination, and the severity of the particular disease being treated and form of administration.

The compounds can be administered with other neurotrophic agents such as neurotrophic growth factor (NGF), glial derived growth factor, brain derived growth factor, ciliary neurotrophic factor, and neurotrophin-3. The dosage level of other neurotrophic drugs will depend upon the factors previously stated and the neurotrophic effectiveness of the drug combination.

As discussed above, the compounds of the present invention have an affinity for the FK506 binding protein, particularly FKBP12. The inhibition of the prolyl peptidyl cis-trans isomerase activity of FKBP may be measured as an indicator of this affinity.

#### Ki Test Procedure

Inhibition of the peptidyl-prolyl isomerase (rotamase) activity of the inventive compounds can be evaluated by known methods described in the literature (Harding, et al., Nature, 1989, 341:758-760; Holt et al. J. Am. Chem. Soc., 115:9923-9938). These values

as apparent  $K_i$ 's and are presented in Table II. The *cis-trans* isomerization of an alanine-proline bond in a model substrate, N-succinyl-Ala-Ala-Pro-Phe-p-nitroanilide, is monitored spectrophotometrically in a chymotrypsin-coupled assay, which releases para-nitroanilide from the *trans* form of the substrate. The inhibition of this reaction caused by the addition of different concentrations of inhibitor is determined, and the data is analyzed as a change in first-order rate constant as a function of inhibitor concentration to yield the apparent  $K_i$  values.

In a plastic cuvette are added 950  $\mu$ L of ice cold assay buffer (25 mM HEPES, pH 7.8, 100 mM NaCl), 10  $\mu$ L of FKBP (2.5 mM dithiothreitol), 25  $\mu$ L of chymotrypsin (50 mg/ml in 1 mM HCL) and 10  $\mu$ L of test compound at various concentrations in dimethyl sulfoxide. The reaction is initiated by the addition of 5  $\mu$ L of substrate (succinyl-Ala-Phe-Pro-Phe-para-nitroanilide, 5 mg/mL in 2.35 mM LiCl in trifluoroethanol).

The absorbance at 390 nm versus time is monitored for 90 seconds using a spectrophotometer and the rate constants are determined from the absorbance versus time data files.

The data for these experiments for representative compounds are presented in Table II under the column " $K_i$ ".

The neurotrophic effects of the compounds of the

present invention can be demonstrated in cellular biological experiments in vitro, as described below.

### Chick Dorsal Root Ganglion

5

### Cultures and Neurite Growth

Dorsal root ganglia were dissected from chick embryos of ten day gestation. Whole ganglion explants were cultured on thin layer Matrigel-coated 12 well plates with Liebovitz L15 plus high glucose media supplemented with 2 mM glutamine and 10% fetal calf serum, and also containing 10  $\mu$ M cytosine  $\beta$ -D arabinofuranoside (Ara C) at 37°C in an environment containing 5% CO<sub>2</sub>. Twenty-four hours later, the DRGs were treated with various immunophilin ligands. Forty-eight hours after drug treatment, the ganglia were visualized under phase contrast or Hoffman Modulation contrast with a Zeiss Axiovert inverted microscope. Photomicrographs of the explants were made, and neurite outgrowth was quantified. Neurites longer than the DRG diameter were counted as positive, with total number of neurites quantitated per each experimental condition. Three to four DRGs are cultured per well, and each treatment was performed in duplicate.

25

The unexpectedly greater metabolic stability of the N-oxides of the present invention may be demonstrated in vitro assays. In studies to directly asses the rate of metabolism of N-oxides, a mouse



### Esterase Activity

Purified rabbit liver esterase was purchased from Sigma. Five (5) units of enzyme were placed in 2 mL of 0.05 M Tris buffer (pH 7.5). Incubations were performed for 2 hours. The reaction was terminated by removing an aliquot from the reaction mixture and placing it in a tube with an equal volume of acetonitrile and the bioanalytical internal standard. The results of these experiments are presented in Table II as rates of enzymatic degradation of the compounds.

The results of these *in vitro* experiments for 3-(3-Pyridyl)-1-propyl (2*S*)-1-(3,3-Dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate ("Parent") are tabulated in Table II.

Table II

Compound	$K_i$ (nM)	$ED_{50}$ (nM)	$t_{1/2}$ (min)	esterase degradation (pmol/min/mg/ protein)
Parent	7.5	0.05	8.1	7521
Example 1	225	2.3	42.8	367

### MPTP Model of Parkinson's Disease

The remarkable neurotrophic and neuroregenerative effects of the present inventive compounds can be further demonstrated in an animal model of neurodegenerative disease. MPTP lesioning of

white mice are dosed i.p. with 30 mg/kg of MPTP for 5 days. Test compounds (4 mg/kg), or vehicle, are administered s.c. along with the MPTP for 5 days, as well as for an additional five days following cessation of MPTP treatment. At 18 days following MPTP treatment, the animals are sacrificed and the striata dissected and perfusion-fixed. Immunostaining is performed on sagittal and coronal brain sections using anti-tyrosine hydroxylase 1 g to quantitative survival and recovery of dopaminergic neurons. In animals treated MPTP and vehicle, a substantial loss of functional dopaminergic terminals is observed as compared to non-lesioned animals. Lesioned animals receiving test compounds show a significant recovery of TH-stained dopaminergic neurons. This model presents quantitation for the recovery of TH-positive dopaminergic neurons in the striatum of animals receiving of animals receiving the compounds of the present invention. Data for representative control and lesioned animals not receiving the test drugs is also presented against the data from the animals receiving the compounds of the present invention.

Data for the representative control and lesioned animals not receiving test drugs is presented against the Parent compound and Example 1 of the present invention in Figure 1. It is clear that, although

Example 1 is less potent in vitro, it is more potent in this in vivo model of neurodegeneration, due to its unexpectedly better bioavailability and pharmacokinetics.

5

### Examples

The following examples are illustrative of the present invention and are not intended to be limitations thereon.

10

### Example 1

Synthesis of 3-(3-Pyridyl)-1-propyl (2S)-1-(3,3-Dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate, N-oxide. (1)

15

Methyl (2S)-1-(1,2-dioxo-2-methoxyethyl)-2-pyrrolidinecarboxylate. A solution of L-proline methyl ester hydrochloride (3.08 g; 18.60 mmol) in dry methyl chloride was cooled to 0°C and treated with triethylamine (3.92 g; 38.74 mmol; 2.1 eq). After stirring the formed slurry under a nitrogen atmosphere for 15 min, a solution of methyl oxalyl chloride (3.20 g; 26.12 mmol) in methylene chloride (45 mL) was added dropwise. The resulting mixture was stirred at 0°C for 1.5 hr. After filtering to remove solids, the organic phase was washed with water, dried over MgSO<sub>4</sub> and concentrated. The crude residue was purified on a silica gel column, eluting with 50% ethyl acetate in

20

25

hexane, to obtain 3.52 g (88%) of the product as a reddish oil. Mixture of *cis-trans* amide rotamers; data for *trans* rotamer given.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  1.93 (dm, 2H); 2.17 (m, 2H); 3.62 (m, 2H); 3.71 (s, 3H); 3.79, 3.84 (s, 3H total); 4.86 (dd, 1H,  $J=8.4, 3.3$ ).

Methyl (2*S*)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidinecarboxylate. A solution of methyl (2*S*)-1-(1,2-dioxo-2-methoxyethyl)-2-pyrrolidinecarboxylate (2.35 g; 10.90 mmol) in 30 mL of tetrahydrofuran (THF) was cooled to  $-78^\circ\text{C}$  and treated with 14.2 mL of a 1.0 M solution of 1,1-dimethylpropylmagnesium chloride in THF. After stirring the resulting homogeneous mixture at  $-78^\circ\text{C}$  for three hours, the mixture was poured into saturated ammonium chloride (100 mL) and extracted into ethyl acetate. The organic phase was washed with water, dried, and concentrated, and the crude material obtained upon removal of the solvent was purified on a silica gel column, eluting with 25% ethyl acetate in hexane, to obtain 2.10 g (75%) of the oxamate as a colorless oil.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  0.88 (t, 3H); 1.22, 1.26 (s, 3H each); 1.75 (dm, 2H); 1.87-2.10 (m, 3H); 2.23 (m, 1H); 3.54 (m, 2H); 3.76 (s, 3H); 4.52 (dm, 1H,  $J=8.4, 3.4$ ).

(2*S*)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidinecarboxylic acid. A mixture of methyl (2*S*)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-



pyrrolidinecarboxylate (2.10 g; 8.23 mmol), 1 N LiOH (15 mL), and methanol (50 mL) was stirred at 0°C for 30 min and at room temperature overnight. The mixture was acidified to pH 1 with 1 N HCl, diluted with water, and extracted into 100 mL of methylene chloride. The organic extract was washed with brine and concentrated to deliver 1.73 g (87%) of snow-white solid which did not require further purification. <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 0.87 (t, 3H); 1.22, 1.25 (s, 3H each); 1.77 (dm, 2H); 2.02 (m, 2H); 2.17 (m, 1H); 2.25 (m, 1H); 3.53 (dd, 2H, J=10.4, 7.3); 4.55 (dd, 1H, J=8.6, 4.1).


3-(3-Pyridyl)-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate. A mixture of (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidinecarboxylic acid (4.58 g; 19 mmol), 3-pyridinepropanol (3.91 g; 28.5 mmol), dicyclohexylcarbodiimide (6.27 g; 30.4 mmol), camphorsulphonic acid (1.47 g; 6.33 mmol) and 4-dimethyl aminopyridine (773 mg; 6.33 mmol) in methylene chloride (100 mL) was stirred overnight under a nitrogen atmosphere. The reaction mixture was filtered through Celite to remove solids and concentrate in vacuo. The crude material was triturated with several portions of ether, and the ether portions were filtered through Celite to remove solids and concentrated in vacuo. The concentrated filtrate was purified on a flash column (gradient

elution, 25% ethyl acetate in hexane to pure ethyl acetate) to obtain 5.47 g (80%) of GPI 1046 as a colorless oil (partial hydrate).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 300 MHz):  $\delta$  0.85 (t, 3H); 1.23, 1.26 (s, 3H each); 1.63-1.89 (m, 2H); 1.90-2.30 (m, 4H); 2.30-2.50 (m, 1H); 2.72 (t, 2H); 3.53 (m, 2H); 4.19 (m, 2H); 4.53 (m, 1H); 7.22 (m, 1H); 7.53 (dd, 1H); 8.45. Anal. Calcd. for  $\text{C}_{20}\text{H}_{28}\text{NO}_4 \cdot 0.25 \text{H}_2\text{O}$ : C, 65.82; H, 7.87; N, 7.68. Found: C, 66.01; H, 7.85; N, 7.64.

3-(3-Pyridyl)-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate, N-oxide. A solution of 3-(3-pyridyl)-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate (190 mg; 0.52 mmol) and m-chloroperbenzoic acid (160 mg of 57%-86% material, 0.53 mmol) was stirred in methylene chloride (20mL) at room temperature for 3 hrs. The reaction mixture was diluted with methylene chloride and washed twice with 1N NaOH. The organic extract was dried and concentrated and the crude material was chromatographed, eluting with 10% methanol in ethyl acetate, to obtain 130 mg of the compound of Example 1,  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 300 MHz):  $\delta$  0.83 (t, 3H); 1.21 (s, 3H); 1.25 (s, 3H); 1.75-2.23 (m, 8H); 2.69 (t, 2H, J=7.5); 3.52 (t, 2H, J=6.3); 4.17 (dd, 2H, J=6.3); 4.51 (m, 1H); 7.16-7.22 (m, 2H); 8.06-8.11 (m, 2H). Anal. Calcd. for  $\text{C}_{20}\text{H}_{28}\text{N}_2\text{O}_5 \cdot 0.75 \text{H}_2\text{O}$ : C, 61.60; H, 7.63; N, 7.18. Found: C, 61.79; H, 7.58; N, 7.23.

## WHAT IS CLAIMED IS:

1. A compound of the formula



or a pharmaceutically acceptable salt thereof,  
 5 wherein:

A and B are taken together, with the nitrogen and carbon atoms to which they are respectively attached, to form a 5-7 membered saturated or unsaturated heterocyclic ring containing any combination of CH<sub>2</sub>, O, S, SO, SO<sub>2</sub>, NH or NR<sub>1</sub> in any chemically stable  
 10 oxidation state;

W is O, S, CH<sub>2</sub>, or H<sub>2</sub>;

R is a C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or alkenyl group optionally substituted with C<sub>3</sub>-C<sub>8</sub> cycloalkyl, C<sub>3</sub> or C<sub>5</sub> cycloalkyl, C<sub>5</sub>-C<sub>7</sub> cycloalkenyl, or Ar<sub>1</sub>, where said alkyl, alkenyl, cycloalkyl, or cycloalkenyl groups may be optionally substituted with C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkenyl, or hydroxy, and where Ar<sub>1</sub> is  
 15

selected from the group consisting of 1-naphthyl, 2-naphthyl, 1-indolyl, 2-indolyl, 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, or phenyl, having one to three substituents which are  
5 independently selected from the group consisting of hydrogen, halo, hydroxyl, nitro, trifluoromethyl, C<sub>1</sub>-C<sub>6</sub> straight or branched alkyl or alkenyl, C<sub>1</sub>-C<sub>4</sub> alkenyloxy, phenoxy, benzyloxy, and amino;

X is O, NH, NR<sub>1</sub>, S, CH, CR<sub>1</sub>, or C(R<sub>1</sub>)<sub>2</sub>;

10 Y is a direct bond, or a C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or alkenyl which is optionally substituted in one or more positions with C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or alkenyl, or C<sub>3</sub>-C<sub>9</sub> cycloalkyl, or C<sub>5</sub>-C<sub>7</sub> cycloalkenyl, or hydroxyl, or  
15 carbonyl oxygen, or with Ar, where said alkyl, alkenyl, cycloalkyl, cycloalkenyl, or Ar group is optionally substituted with C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkenyl, or hydroxy, or carbonyl oxygen, or wherein any of the carbon atoms of said alkyl, alkenyl, cycloalkyl,  
20 cycloalkenyl, or Ar group is optionally replaced with O, NH, NR<sub>2</sub>, S, SO, or SO<sub>2</sub>, where R<sub>2</sub> is selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>4</sub>)-straight or branched chain alkyl, (C<sub>3</sub>-C<sub>4</sub>)-straight or branched chain alkenyl or alkynyl, and (C<sub>1</sub>-C<sub>4</sub>) bridging alkyl  
25 wherein a bridge is formed between the nitrogen and a carbon atom of said alkyl or alkenyl chain containing said heteroatom to form a ring, wherein said ring is optionally fused to an Ar group; and

Z is an aromatic or tertiary alkyl amine oxidized to a corresponding N-oxide, wherein the aromatic amine is Ar oxidized to a corresponding N-oxide where Ar is a mono-, bi- or tricyclic, carbo- or heterocyclic ring, wherein the ring is either unsubstituted or substituted in one to three position(s) with halo, hydroxyl, nitro, trifluoromethyl, C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or alkenyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> alkenyloxy, phenoxy, benzyloxy, amino, or a combination thereof; wherein the individual ring sizes are 5-6 members; wherein the heterocyclic ring contains 1-6 heteroatom(s) selected from the group consisting of O, N, S, and a combination thereof wherein at least one of the heteroatoms is N, and wherein the alkyl amine is oxidized to a corresponding N-oxide where alkyl is a C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or alkenyl which is optionally substituted in one or more positions with C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or alkenyl, or C<sub>3</sub>-C<sub>8</sub> cycloalkyl, or C<sub>5</sub>-C<sub>7</sub> cycloalkenyl, or hydroxyl, or carbonyl oxygen, or with Ar, where said alkyl, alkenyl, cycloalkyl, cycloalkenyl, or Ar group is optionally substituted with C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkenyl, or hydroxy, or carbonyl oxygen, or wherein any of the carbon atoms of said alkyl, alkenyl, cycloalkyl, cycloalkenyl, or Ar group is optionally replaced with O, NH, NR<sub>1</sub>, S, SO, or SO<sub>2</sub>;

Ar is selected from the group consisting of pyrrolidinyl, pyridyl, pyrimidyl, pyrazyl, pyridazyl,

quinolinyl, and isoquinolinyl; and,

$R_1$  is hydrogen,  $(C_1-C_4)$ -straight or branched chain alkyl,  $(C_3-C_4)$ -straight or branched chain alkenyl or alkynyl, or  $R_1$  is Y-Z as defined above.

5

2. The compound of claim 1, which has an affinity for FKBP type immunophilins.

10

3. The compound of claim 2, where the FKBP-type immunophilin is FKBP12.

4. The compound of claim 1, capable of inhibiting rotamase enzyme activity.

15

5. A pharmaceutical composition comprising a neurotrophically effective amount of the compound of claim 1 and a pharmaceutically acceptable carrier.

20

6. A method of stimulating damaged neurons in an animal, comprising:

administering to the animal a neurotrophically effective amount of the compound of claim 1.

25

7. A method of promoting neuronal regeneration in an animal, comprising:

administering to the animal a neurotrophically effective amount of the compound of claim 1.

8. A method of preventing neurodegeneration in an animal, comprising:

administering to the animal a neurotrophically effective amount of the compound of claim 1.

5

9. A method of treating neurological disorders in an animal, comprising:

administering to the animal a neurotrophically effective amount of the compound of claim 1.

10


10. The method of claim 9, wherein the neurological disorder is selected from the group consisting of peripheral neuropathy caused by physical injury or disease state, physical damage to the brain, physical damage to the spinal cord, stroke associated with brain damage, and neurological disorder relating to neurodegeneration.

15

11. The method of claim 10, wherein the neurological disorder relating to neurodegeneration is selected from the group consisting of Alzheimer's Disease, Parkinson's Disease, and amyotrophic lateral sclerosis.

20

12. A compound of formula II comprising:



II

5 or a pharmaceutically acceptable salt thereof,  
wherein:

E, F, G and H are independently  $\text{CH}_2$ , O, S, SO,  $\text{SO}_2$ ,  
NH or  $\text{NR}_1$ ;

W is O, S,  $\text{CH}_2$ , or  $\text{H}_2$ ;

10 R is a  $\text{C}_1$ - $\text{C}_6$  straight or branched chain alkyl or  
alkenyl group optionally substituted with  $\text{C}_3$ - $\text{C}_8$   
cycloalkyl,  $\text{C}_3$  or  $\text{C}_5$  cycloalkyl,  $\text{C}_5$ - $\text{C}_7$  cycloalkenyl, or  
 $\text{Ar}_1$ , where said alkyl, alkenyl, cycloalkyl, or  
cycloalkenyl groups may be optionally substituted with  
15  $\text{C}_1$ - $\text{C}_4$  alkyl,  $\text{C}_1$ - $\text{C}_4$  alkenyl, or hydroxy, and where  $\text{Ar}_1$  is  
selected from the group consisting of 1-naphthyl, 2-  
naphthyl, 1-indolyl, 2-indolyl, 2-furyl, 3-furyl, 2-  
thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl,  
or phenyl, having one to three substituents which are  
20 independently selected from the group consisting of  
hydrogen, halo, hydroxyl, nitro, trifluoromethyl,  $\text{C}_1$ - $\text{C}_6$   
straight or branched alkyl or alkenyl,  $\text{C}_1$ - $\text{C}_4$   
alkenyloxy, phenoxy, benzyloxy, and amino;

X is O, NH,  $\text{NR}_1$ , S, CH,  $\text{CR}_1$ , or  $\text{C}(\text{R}_1)_2$ ;



Y is a direct bond, or a C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or alkenyl which is optionally substituted in one or more positions with C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or alkenyl, or C<sub>3</sub>-C<sub>8</sub> cycloalkyl, or C<sub>5</sub>-C<sub>7</sub> cycloalkenyl, or hydroxyl, or carbonyl oxygen, or with Ar, where said alkyl, alkenyl, cycloalkyl, cycloalkenyl, or Ar group is optionally substituted with C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkenyl, or hydroxy, or carbonyl oxygen, or wherein any of the carbon atoms of said alkyl, alkenyl, cycloalkyl, cycloalkenyl, or Ar group is optionally replaced with O, NH, NR<sub>2</sub>, S, SO, or SO<sub>2</sub>, where R<sub>2</sub> is selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>4</sub>)-straight or branched chain alkyl, (C<sub>3</sub>-C<sub>4</sub>)-straight or branched chain alkenyl or alkynyl, and (C<sub>1</sub>-C<sub>4</sub>) bridging alkyl wherein a bridge is formed between the nitrogen and a carbon atom of said alkyl or alkenyl chain containing said heteroatom to form a ring, wherein said ring is optionally fused to an Ar group; and

Z is an aromatic or tertiary alkyl amine oxidized to a corresponding N-oxide, wherein the aromatic amine is Ar oxidized to a corresponding N-oxide where Ar is a mono-, bi- or tricyclic, carbo- or heterocyclic ring, wherein the ring is either unsubstituted or substituted in one to three position(s) with halo, hydroxyl, nitro, trifluoromethyl, C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or alkenyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> alkenyloxy, phenoxy, benzyloxy, amino, or a

combination thereof; wherein the individual ring sizes are 5-6 members; wherein the heterocyclic ring contains 1-6 heteroatom(s) selected from the group consisting of O, N, S, and a combination thereof wherein at least one of the heteroatoms is N, and wherein the alkyl amine is oxidized to a corresponding N-oxide where alkyl is a C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or alkenyl which is optionally substituted in one or more positions with C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or alkenyl, or C<sub>3</sub>-C<sub>8</sub> cycloalkyl, or C<sub>5</sub>-C<sub>7</sub> cycloalkenyl, or hydroxyl, or carbonyl oxygen, or with Ar, where said alkyl, alkenyl, cycloalkyl, cycloalkenyl, or Ar group is optionally substituted with C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkenyl, or hydroxy, or carbonyl oxygen, or wherein any of the carbon atoms of said alkyl, alkenyl, cycloalkyl, cycloalkenyl, or Ar group is optionally replaced with O, NH, NR<sub>1</sub>, S, SO, or SO<sub>2</sub>;

Ar is selected from the group consisting of pyrrolidinyl, pyridyl, pyrimidyl, pyrazyl, pyridazyl, quinolinyl, and isoquinolinyl; and,

R<sub>1</sub> is hydrogen, (C<sub>1</sub>-C<sub>4</sub>)-straight or branched chain alkyl, (C<sub>3</sub>-C<sub>4</sub>)-straight or branched chain alkenyl or alkynyl, or R<sub>1</sub> is Y-Z as defined above.

13. The compound of claim 12, which has an affinity for FKBP type immunophilins.

14. The compound of claim 13, where the FKBP-type

immunophilin is FKBP12.

15. The compound of claim 12, capable of inhibiting rotamase enzyme activity.

5

16. A pharmaceutical composition comprising a neurotrophically effective amount of the compound of claim 12 and a pharmaceutically acceptable carrier.

10 17. A method of stimulating damaged neurons in an animal, comprising:

administering to the animal a neurotrophically effective amount of the compound of claim 12.

15 18. A method of promoting neuronal regeneration in an animal, comprising:

administering to the animal a neurotrophically effective amount of the compound of claim 12.

20 19. A method of preventing neurodegeneration in an animal, comprising:

administering to the animal a neurotrophically effective amount of the compound of claim 12.


25 20. A method of treating neurological disorders in an animal, comprising:

administering to the animal a neurotrophically effective amount of the compound of claim 12.

brain damage, and neurological disorder relating to neurodegeneration.

22. The method of claim 21, wherein the neurological  
10 disorder relating to neurodegeneration is selected  
from the group consisting of Alzheimer's Disease,  
Parkinson's Disease, and amyotrophic lateral  
sclerosis.

23. A compound of formula III comprising:



III

5 or a pharmaceutically acceptable salt thereof,  
wherein:

E, F, and G are independently  $\text{CH}_2$ , O, S,  $\text{SO}$ ,  $\text{SO}_2$ ,  
NH or  $\text{NR}_1$ ;

W is O, S,  $\text{CH}_2$ , or  $\text{H}_2$ ;

10 R is a  $\text{C}_1$ - $\text{C}_6$  straight or branched chain alkyl or  
alkenyl group optionally substituted with  $\text{C}_3$ - $\text{C}_8$   
cycloalkyl,  $\text{C}_3$  or  $\text{C}_5$  cycloalkyl,  $\text{C}_5$ - $\text{C}_7$  cycloalkenyl, or  
 $\text{Ar}_1$ , where said alkyl, alkenyl, cycloalkyl, or  
cycloalkenyl groups may be optionally substituted with  
15  $\text{C}_1$ - $\text{C}_4$  alkyl,  $\text{C}_1$ - $\text{C}_4$  alkenyl, or hydroxy, and where  $\text{Ar}_1$  is  
selected from the group consisting of 1-naphthyl, 2-  
naphthyl, 1-indolyl, 2-indolyl, 2-furyl, 3-furyl, 2-  
thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl,  
or phenyl, having one to three substituents which are  
20 independently selected from the group consisting of  
hydrogen, halo, hydroxyl, nitro, trifluoromethyl,  $\text{C}_1$ - $\text{C}_6$   
straight or branched alkyl or alkenyl,  $\text{C}_1$ - $\text{C}_4$   
alkenyloxy, phenoxy, benzyloxy, and amino;

X is O, NH,  $\text{NR}_1$ , S, CH,  $\text{CR}_1$ , or  $\text{C}(\text{R}_1)_2$ ;

Y is a direct bond, or a C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or alkenyl which is optionally substituted in one or more positions with C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or alkenyl, or C<sub>3</sub>-C<sub>8</sub> cycloalkyl, or C<sub>5</sub>-C<sub>7</sub> cycloalkenyl, or hydroxyl, or carbonyl oxygen, or with Ar, where said alkyl, alkenyl, cycloalkyl, cycloalkenyl, or Ar group is optionally substituted with C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkenyl, or hydroxy, or carbonyl oxygen, or wherein any of the carbon atoms of said alkyl, alkenyl, cycloalkyl, cycloalkenyl, or Ar group is optionally replaced with O, NH, NR<sub>2</sub>, S, SO, or SO<sub>2</sub>, where R<sub>2</sub> is selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>4</sub>)-straight or branched chain alkyl, (C<sub>3</sub>-C<sub>4</sub>)-straight or branched chain alkenyl or alkynyl, and (C<sub>1</sub>-C<sub>4</sub>) bridging alkyl wherein a bridge is formed between the nitrogen and a carbon atom of said alkyl or alkenyl chain containing said heteroatom to form a ring, wherein said ring is optionally fused to an Ar group; and

Z is an aromatic or tertiary alkyl amine oxidized to a corresponding N-oxide, wherein the aromatic amine is Ar oxidized to a corresponding N-oxide where Ar is a mono-, bi- or tricyclic, carbo- or heterocyclic ring, wherein the ring is either unsubstituted or substituted in one to three position(s) with halo, hydroxyl, nitro, trifluoromethyl, C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or alkenyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> alkenyloxy, phenoxy, benzyloxy, amino, or a

combination thereof; wherein the individual ring sizes are 5-6 members; wherein the heterocyclic ring contains 1-6 heteroatom(s) selected from the group consisting of O, N, S, and a combination thereof wherein at least one of the heteroatoms is N, and wherein the alkyl amine is oxidized to a corresponding N-oxide where alkyl is a C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or alkenyl which is optionally substituted in one or more positions with C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or alkenyl, or C<sub>3</sub>-C<sub>8</sub> cycloalkyl, or C<sub>5</sub>-C<sub>7</sub> cycloalkenyl, or hydroxyl, or carbonyl oxygen, or with Ar, where said alkyl, alkenyl, cycloalkyl, cycloalkenyl, or Ar group is optionally substituted with C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkenyl, or hydroxy, or carbonyl oxygen, or wherein any of the carbon atoms of said alkyl, alkenyl, cycloalkyl, cycloalkenyl, or Ar group is optionally replaced with O, NH, NR<sub>1</sub>, S, SO, or SO<sub>2</sub>;

Ar is selected from the group consisting of pyrrolidinyl, pyridyl, pyrimidyl, pyrazyl, pyridazyl, quinolinyl, and isoquinolinyl; and,

R<sub>1</sub> is hydrogen, (C<sub>1</sub>-C<sub>4</sub>)-straight or branched chain alkyl, (C<sub>3</sub>-C<sub>4</sub>)-straight or branched chain alkenyl or alkynyl, or R<sub>1</sub> is Y-Z as defined above.

24. The compound of claim 23, which has an affinity for FKBP type immunophilins.

25. The compound of claim 24, where the FKBP-type

immunophilin is FKBP12.

26. The compound of claim 23, capable of inhibiting rotamase enzyme activity.

5

27. A pharmaceutical composition comprising a neurotrophically effective amount of the compound of claim 23 and a pharmaceutically acceptable carrier.

10 28. A method of stimulating damaged neurons in an animal, comprising:

administering to the animal a neurotrophically effective amount of the compound of claim 23.

15 29. A method of promoting neuronal regeneration in an animal, comprising:

administering to the animal a neurotrophically effective amount of the compound of claim 23.

20 30. A method of preventing neurodegeneration in an animal, comprising:

administering to the animal a neurotrophically effective amount of the compound of claim 23.

25 31. A method of treating neurological disorders in an animal, comprising:


administering to the animal a neurotrophically effective amount of the compound of claim 23.



32. The method of claim 31, wherein the neurological disorder is selected from the group consisting of peripheral neuropathy caused by physical injury or disease state, physical damage to the brain, physical damage to the spinal cord, stroke associated with brain damage, and neurological disorder relating to neurodegeneration.

33. The method of claim 32, wherein the neurological disorder relating to neurodegeneration is selected from the group consisting of Alzheimer's Disease, Parkinson's Disease, and amyotrophic lateral sclerosis.

34. A compound of formula IV comprising:



IV

or a pharmaceutically acceptable salt thereof, wherein:

$n$  is 1, 2 or 3 forming a 5-7 member heterocyclic ring;

$W$  is O, S,  $CH_2$ , or  $H_2$ ;

$R$  is a  $C_1$ - $C_6$  straight or branched chain alkyl or alkenyl group optionally substituted with  $C_3$ - $C_8$

cycloalkyl, C<sub>3</sub> or C<sub>5</sub> cycloalkyl, C<sub>3</sub>-C<sub>7</sub> cycloalkenyl, or Ar<sub>1</sub>, where said alkyl, alkenyl, cycloalkyl, or cycloalkenyl groups may be optionally substituted with C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkenyl, or hydroxy, and where Ar<sub>1</sub> is  
5 selected from the group consisting of 1-naphthyl, 2-naphthyl, 1-indolyl, 2-indolyl, 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, or phenyl, having one to three substituents which are independently selected from the group consisting of  
10 hydrogen, halo, hydroxyl, nitro, trifluoromethyl, C<sub>1</sub>-C<sub>6</sub> straight or branched alkyl or alkenyl, C<sub>1</sub>-C<sub>4</sub> alkenyloxy, phenoxy, benzyloxy, and amino;

X is O, NH, NR<sub>1</sub>, S, CH, CR<sub>1</sub>, or C(R<sub>1</sub>)<sub>2</sub>;

Y is a direct bond, or a C<sub>1</sub>-C<sub>6</sub> straight or  
15 branched chain alkyl or alkenyl which is optionally substituted in one or more positions with C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or alkenyl, or C<sub>3</sub>-C<sub>8</sub> cycloalkyl, or C<sub>5</sub>-C<sub>7</sub> cycloalkenyl, or hydroxyl, or carbonyl oxygen, or with Ar, where said alkyl, alkenyl, cycloalkyl, cycloalkenyl, or Ar group is  
20 optionally substituted with C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkenyl, or hydroxy, or carbonyl oxygen, or wherein any of the carbon atoms of said alkyl, alkenyl, cycloalkyl, cycloalkenyl, or Ar group is optionally replaced with  
25 O, NH, NR<sub>2</sub>, S, SO, or SO<sub>2</sub>, where R<sub>2</sub> is selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>4</sub>)-straight or branched chain alkyl, (C<sub>3</sub>-C<sub>4</sub>)-straight or branched chain alkenyl or alkynyl, and (C<sub>1</sub>-C<sub>4</sub>) bridging alkyl

wherein a bridge is formed between the nitrogen and a carbon atom of said alkyl or alkenyl chain containing said heteroatom to form a ring, wherein said ring is optionally fused to an Ar group; and

5           Z is an aromatic or tertiary alkyl amine oxidized to a corresponding N-oxide, wherein the aromatic amine is Ar oxidized to a corresponding N-oxide where Ar is a mono-, bi- or tricyclic, carbo- or heterocyclic ring, wherein the ring is either unsubstituted or  
10 substituted in one to three position(s) with halo, hydroxyl, nitro, trifluoromethyl, C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or alkenyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> alkenyloxy, phenoxy, benzyloxy, amino, or a combination thereof; wherein the individual ring sizes  
15 are 5-6 members; wherein the heterocyclic ring contains 1-6 heteroatom(s) selected from the group consisting of O, N, S, and a combination thereof wherein at least one of the heteroatoms is N, and wherein the alkyl amine is oxidized to a corresponding  
20 N-oxide where alkyl is a C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or alkenyl which is optionally substituted in one or more positions with C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or alkenyl, or C<sub>3</sub>-C<sub>8</sub> cycloalkyl, or C<sub>5</sub>-C<sub>8</sub> cycloalkenyl, or hydroxyl, or carbonyl oxygen,  
25 or with Ar, where said alkyl, alkenyl, cycloalkyl, cycloalkenyl, or Ar group is optionally substituted with C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkenyl, or hydroxy, or carbonyl oxygen, or wherein any of the carbon atoms of said

alkyl, alkenyl, cycloalkyl, cycloalkenyl, or Ar group is optionally replaced with O, NH, NR<sub>1</sub>, S, SO, or SO<sub>2</sub>;

Ar is selected from the group consisting of pyrrolidinyl, pyridyl, pyrimidyl, pyrazyl, pyridazyl, quinolinyl, and isoquinolinyl; and,

R<sub>1</sub> is hydrogen, (C<sub>1</sub>-C<sub>4</sub>)-straight or branched chain alkyl, (C<sub>3</sub>-C<sub>4</sub>)-straight or branched chain alkenyl or alkynyl, or R<sub>1</sub> is Y-Z as defined above.

35. The compound of claim 34, which has an affinity for FKBP type immunophilins.

36. The compound of claim 35, where the FKBP-type immunophilin is FKBP12.

37. The compound of claim 34, capable of inhibiting rotamase enzyme activity.

38. A pharmaceutical composition comprising a neurotrophically effective amount of the compound of claim 34 and a pharmaceutically acceptable carrier.

39. A method of stimulating damaged neurons in an animal, comprising:

administering to the animal a neurotrophically effective amount of the compound of claim 34.

40. A method of promoting neuronal regeneration in an

animal, comprising:

administering to the animal a neurotrophically effective amount of the compound of claim 34.

- 5 41. A method of preventing neurodegeneration in an animal, comprising:

administering to the animal a neurotrophically effective amount of the compound of claim 34.

- 10 42. A method of treating neurological disorders in an animal, comprising:

administering to the animal a neurotrophically effective amount of the compound of claim 34.

- 15 43. The method of claim 42, wherein the neurological disorder is selected from the group consisting of peripheral neuropathy caused by physical injury or disease state, physical damage to the brain, physical damage to the spinal cord, stroke associated with  
20 brain damage, and neurological disorder relating to neurodegeneration.

44. The method of claim 43, wherein the neurological disorder relating to neurodegeneration is selected  
25 from the group consisting of Alzheimer's Disease, Parkinson's Disease, and amyotrophic lateral sclerosis.

45. A compound selected from the group consisting of:
- 3-(2-Pyridyl)-1-propyl (2S)-1-(1,1-Dimethyl-1,2-dioxo-pentyl)-2-pyrrolidinecarboxylate, N-oxide;
- 3-(3-Pyridyl)-1-propyl (2S)-1-(1,1-Dimethyl-1,2-dioxo-pentyl)-2-pyrrolidinecarboxylate, N-oxide;
- 3-(4-Pyridyl)-1-propyl (2S)-1-(1,1-Dimethyl-1,2-dioxo-pentyl)-2-pyrrolidinecarboxylate, N-oxide;
- 3-(2-Quinolyl)-1-propyl (2S)-1-(1,1-Dimethyl-1,2-dioxo-pentyl)-2-pyrrolidinecarboxylate, N-oxide;
- 3-(3-Quinolyl)-1-propyl (2S)-1-(1,1-Dimethyl-1,2-dioxo-pentyl)-2-pyrrolidinecarboxylate, N-oxide; and
- 3-(4-Quinolyl)-1-propyl (2S)-1-(1,1-Dimethyl-1,2-dioxo-pentyl)-2-pyrrolidinecarboxylate, N-oxide.
46. The compound of claim 45, which has an affinity for FKBP type immunophilins.
47. The compound of claim 46, where the FKBP-type immunophilin is FKBP12.
48. The compound of claim 45, capable of inhibiting rotamase enzyme activity.
49. A pharmaceutical composition comprising a neurotrophically effective amount of the compound of claim 45 and a pharmaceutically acceptable carrier.
50. A method of stimulating damaged neurons in an

animal, comprising:

administering to the animal a neurotrophically effective amount of the compound of claim 45.

5 51. A method of promoting neuronal regeneration in an animal, comprising:

administering to the animal a neurotrophically effective amount of the compound of claim 45.

10 52. A method of preventing neurodegeneration in an animal, comprising:

administering to the animal a neurotrophically effective amount of the compound of claim 45.

15 53. A method of treating neurological disorders in an animal, comprising:

administering to the animal a neurotrophically effective amount of the compound of claim 45.


20 54. The method of claim 53, wherein the neurological disorder is selected from the group consisting of peripheral neuropathy caused by physical injury or disease state, physical damage to the brain, physical damage to the spinal cord, stroke associated with  
25 brain damage, and neurological disorder relating to neurodegeneration.

55. The method of claim 54, wherein the neurological

disorder relating to neurodegeneration is selected from the group consisting of Alzheimer's Disease, Parkinson's Disease, and amyotrophic lateral sclerosis.



FIG. 1



**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(6) :A61K 31/40, 31/44, 31/47; C07D 207/08, 207/09, 213/89, 215/60

US CL :514/279.1, 343; 546/174, 314

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/279.1, 343; 546/174, 314

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Registry, Chemical Abstracts

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y,P	Chem. abstr., Vol.126, No.21, issued 1997 (Columbus, OH, USA), page 137, column 2, the abstract Nio. 272710, 'Neurotrophic Immunophilin Ligands Stimulate Structural and Functional Recovery in neurodegenerative Animal Models.' Proc. Natl. Acad. Sci. U.S.A., 1997, 94(5), 2019-2024.	1-55
Y,P	Chem. abstr., Vol.126, No.21, 26 May 1997 (Columbus, OH, USA), page 88, column 1, the abstract Noc. 272378, ARMISTEAD, 'Methods and Compositions for Stimulating Neurite Growth Using Compounds with Affinity for FKBP12 in Combination with Neurotrophic Factors.' ZA 9604852, 1996.	1-55

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*E* earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
*O* document referring to an oral disclosure, use, exhibition or other means	
*P* document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

06 APRIL 1998

Date of mailing of the international search report

17 JUL 1998

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

D. MARGARET M. MACH

Telephone No. (703) 308-1235

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/03484

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Chem. abstr., Vol 122, No.5, 30 January 1995 (Columbus. OH, USA), page 1178, column 2, the abstract No. 55896, ARMISTEAD, '1-(2-oxoacetyl)piperidine-2-carboxylic Acid Derivatives as Multi-drug-resistant Cancer Cell Sensitizers.' WO 9407858, 1994.	1-55
Y	Chem. abstr., Vol.121, No.1, 4 July 1994 (Columbus. OH, USA), page 20, column 1, the abstract No. 224, HOLT, 'Structure-activity Studies of Synthetic FKBP Ligands as Peptidyl-prolyl Isomerase Inhibitors.' Bioorg. Med. Chem. Lett., 1994, 4(2), 315-320.	1-55
Y,P	US 5,696,135 A (STEINER et al.) 09 December 1997, see abstract and claims.	1-55
Y,P	US 5,620,971 A (ARMISTEAD et al) 15 April 1997, see abstract and claims.	1-55

